

# Type 1 Diabetes Among Sardinian Children Is Increasing

The Sardinian diabetes register for children aged 0–14 years (1989–1999)

ANNA CASU, MD<sup>1</sup>  
CRISTIANA PASCUTTO, PHD<sup>2</sup>  
LUISA BERNARDINELLI, PHD<sup>2,3</sup>

THE SARDINIAN IDDM EPIDEMIOLOGY  
STUDY GROUP\*  
MARCO SONGINI, MD<sup>1</sup>

**OBJECTIVE** — The Sardinian type 1 diabetes register represented the basis to determine the most recent trends and the age distribution of type 1 diabetes incidence among Sardinians <15 years of age during 1989–1999. Part of the data (1989–1998) has been already published by the EURODIAB Group with a lower completeness of ascertainment (87%). The geographical distribution of type 1 diabetes risk was also investigated.

**RESEARCH DESIGN AND METHODS** — The new cases of type 1 diabetes in children aged 0–14 years in Sardinia were prospectively registered from 1989 to 1999 according to the EURODIAB ACE criteria. The completeness of ascertainment calculated applying the capture-recapture method was 91%. Standardized incidence rates and 95% CI were calculated assuming the Poisson distribution. Trend of type 1 diabetes incidence was analyzed using the Poisson regression model. Maps of the geographical distribution of type 1 diabetes risk for the whole time period and separately for 1989–1994 and 1995–1999 were produced applying a Bayesian method.

**RESULTS** — A total of 1,214 type 1 diabetic patients were registered yielding to an overall age- and sex-standardized incidence rate of 38.8/100,000 (95% CI 36.7–41.1). There was a male excess with an overall male-to-female ratio of 1.4 (1.3–1.8). The increase of incidence during the 11 years analyzed was statistically significant ( $P = 0.002$ ) with a yearly increasing rate of 2.8% (1.0–4.7). No evidence of an effect of age and sex on this trend has been found. The geographical distribution of type 1 diabetes relative risk (RR) showed that the highest risk areas are located in the southern and central-eastern part of the island and the lowest risk in the northeastern part, even if most of these differences were not statistically significant. This geographical distribution seemed to remain mainly the same between 1989–1994 and 1995–1999.

**CONCLUSIONS** — The homogeneity of diabetes risk and the increase of incidence over the age-groups in the Sardinian population stress the role of an environmental factor uniformly distributed among the genetically high-risk Sardinians.

*Diabetes Care* 27:1623–1629, 2004

It has been claimed that the incidence of type 1 diabetes has increased in many countries over the past 20–30 years (1). This upward trend, first described in

the high-risk populations of Northern Europe, appears to be a global phenomenon, as shown by several studies analyzing >10 years of registration (2–6). Type 1

diabetes incidence has been rising over the last decade by an yearly average of 3.0% worldwide (7) and 3.2% in Europe (8) with a higher rate of increase among children <5 years of age. Earlier data have demonstrated a marked increase in the point prevalence of type 1 diabetes also in Sardinia during the past decades among young men aged 18 years (9). However, that study was not able to investigate the incidence in recent years, nor the risk among women or among different age-groups.

The high incidence of type 1 diabetes and the peculiar genetic characteristics (10) make Sardinia an interesting place to investigate the epidemiology of the disease. According to previous reports, Sardinia had the second highest type 1 diabetes incidence in the world among children aged <15 years (8,11). The risk seemed to be widespread all over the island (11).

The main purpose of the present work was to characterize the temporal trend of the incidence rate of type 1 diabetes in the period 1989–1999 among children <15 years of age living in Sardinia. We also reported sex and age distribution of incidence during the study period. The geographical distribution of type 1 diabetes has also been investigated applying a Bayesian method to smooth the random variation of the estimates due to the small number of events in each municipality. We also analyzed the temporal variation of the geographical distribution of type 1 diabetes risk.

## RESEARCH DESIGN AND METHODS

### Data collection

A population-based and prospective register of type 1 diabetes occurring in children aged <15 years and living in Sardinia has been maintained since 1989 according to the EURODIAB criteria (11,12). Type 1 diabetes was defined on

From the <sup>1</sup>Diabetes Unit, Department of Internal Medicine, Azienda Ospedaliera Brotzu, Via Peretti, Cagliari, Italy; the <sup>2</sup>Dipartimento di Scienze Sanitarie Applicate e Psicocomportamentali, University of Pavia, Pavia, Italy; and the <sup>3</sup>Medical Research Council Biosstatistics Unit, Cambridge, U.K.

Address correspondence and reprint requests to Marco Songini, MD, Diabetes Unit, Azienda Ospedaliera Brotzu, Via Peretti, 09134 Cagliari, Italy. E-mail: marcosongini@aob.it.

Received for publication 5 August 2003 and accepted in revised form 26 March 2004.

\*A complete list of Sardinian IDDM Epidemiology Study Group members can be found in the APPENDIX.

Abbreviations: PP, posterior probability; SIR, standardized incidence rate.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2004 by the American Diabetes Association.

Table 1—Type 1 diabetes incidences by age and sex among Sardinia children &lt;15 years of age (1989–1999)

	SIR 0–14 years (95% CI)*	0–4 years*	5–9 years*	10–14 years*
Total	38.8 (36.7–41.1)†	29.1	42.4	44.4
Boys	46.8 (43.4–50.2)‡	34.1	47.4	55.7
Girls	32.3 (20.4–35.2)‡	24.6	38.0	33.2
Male-to-female ratio	1.4 (1.3–1.8)	1.5	1.3	1.8
Cases (%)	1,214 (100)	256 (21)	430 (35)	528 (44)

\*Incidences per 100,000 person-years; †age and sex SIR; ‡age SIR.

the basis of clinical diagnosis of idiopathic, insulin-dependent diabetes by a doctor, excluding cases occurring secondary to other conditions. The date of onset was taken as the date of first insulin injection. Specific inclusion criteria for this study were date of diagnosis between 1 January 1989 and 31 December 1999 and permanent residency in Sardinia for at least 6 months before the diabetes onset (this excluded children diagnosed in Sardinia by chance). The same methods of data collection have been used since the beginning of the registration.

### Population data

The mid-year estimates of the population obtained from ISTAT (National Institute of Statistics) were available by sex, year of age, calendar year, and municipality for each of the 11 years included in the present investigation (13). The Sardinian population aged <15 years decreased from 317,571 in 1989 to 240,251 in 1999.

### Ascertainment of cases

The completeness of ascertainment was calculated according to the capture-recapture method (14). Diabetes clinics for children and adults and departments of internal medicine and endocrinology were the first source of ascertainment. The second independent source was the membership lists of the local patient associations and the personal National Health System identification cards needed by each patient to obtain medication and devices free of charge.

The first source identified only 790 case subjects, the second source identified 73, and 423 were identified by both sources. The estimated degree of ascertainment was 91% for the whole period investigated.

### Statistical analysis

Total, sex-, and age-specific incidence rates per 100,000 person-years were cal-

culated for each calendar year. A direct standardization method was used to estimate age and sex standardized rates, assuming a reference population comprising equal numbers in each of the sex- and age-specific groups (0–4, 5–9, and 10–14 years). The 95% CI for the standardized incidence rates (SIRs) were calculated using the Gaussian approximation to the Poisson log-likelihood. The time trend of the SIR was assessed using the Poisson regression analysis, including sex and age in the model. This analysis refers to the 5-year age-groups (0–4, 5–9, and 10–14 years) and separately to the 0- to 2-year age-group.

To study the geographical variation of type 1 diabetes risk and filter out the random variation from the maps, we mapped the smoothed estimates of RR at municipality level obtained via a Bayesian model. It is a Poisson Bayesian hierarchical model allowing for spatial correlation of the area-specific RRs. The spatially structured random effects were modeled by means of a conditional autoregressive prior where the mean of each area-specific random effect depends on the random effects of the adjacent areas. As a result, relative risk estimates tend to vary smoothly in space. The model has been fully described previously (15).


The area-specific RR has been calculated as a ratio between the area-specific incidence and the overall incidence in the whole island during the 11 years considered. The Bayesian posterior probability (PP) that the risk in each municipality exceeds the overall incidence was also calculated. This can be interpreted as a “Bayesian *P* value” for the null hypothesis that the area-specific RR is equal to 1. To further evaluate the temporal variations of the geographical distribution of the RR, the periods 1989–1994 and 1995–1999 were analyzed separately to compare the present results with our previous analysis (11).

**RESULTS**— During the 11-year period from 1989 to 1999, a total of 1,214 patients (736 boys, 478 girls) with newly diagnosed type 1 diabetes <15 years of age were recorded. The overall age and sex SIRs in Sardinian children were 38.8/100,000 person-years (95% CI 36.7–41.1).

The incidence rates in the three age-groups (0–4, 5–9, and 10–14 years) by sex are shown in Table 1. The incidence was higher among boys than among girls, yielding an overall male-to-female ratio of 1.4 (95% CI 1.3–1.8). The highest male-to-female ratio was registered in the 10- to 14-year age-group (1.8). The highest incidence rate was registered among boys aged 10–14 years (55.7/100,000). Among girls, the highest incidence was registered in the 5- to 9-year age-group (38.0/100,000). The median age at onset was 9 years for both sexes. The peak of incidence was at 9 years of age for girls, whereas it was at a slightly older age among boys (12 years of age). For both sexes, incidence was quite high at the age of 3 years (41.0/100,000). The incidence rates tend to be higher among boys before the age of 3 years and after the age of 6 years (data not shown).

We observed a rising trend of incidence from 37.7/100,000 at the beginning of the study in 1989 up to 49.3/100,000 at the end in 1999. According to the Poisson regression analysis for a linear time trend adjusted by age and sex, the increasing trend of incidence during the 11 years analyzed was statistically significant ( $P = 0.002$ ) with an estimated average yearly increase of 2.8% (95% CI 1.0–4.7) (Fig. 1). Results by age and sex are presented in Table 2.

To better understand the putative effect of an earlier presentation of the onset, the incidence between 0 and 2 years of age has been analyzed, yielding a lower incidence if the age of onset considered was <1 year (10.3/100,000 [95% CI 6.4–16.4]) and much higher if we considered



Year	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Nr	125	103	103	118	111	91	122	106	114	99	122
Incidence	37.7	33.0	33.2	39.8	38.1	32.0	44.3	39.5	43.5	40.0	49.3
95% C.I.	31.6 45.1	27.1 40.1	27.3 40.3	33.2 47.8	31.6 45.9	26.0 39.3	37.0 52.9	32.6 47.9	36.1 52.3	32.8 48.7	41.3 59.0

**Figure 1**—Incidence rate trend in Sardinia among children aged 0–14 years during 1989–1999 ( $P = 0.002$ ).

the age 1 (29.5/100,000 [22.4–38.9]) and 2 (19.9/100,000 [15.5–25.5]) years. When the temporal trend 0–2 years was analyzed, no significant increase was found for both sexes as well as for girls, whereas an increase was found among boys (7.5% per year [0.3–15.1]). The temporal trend seems to have two peaks: in 1993 and in 1998. This is different from the Finnish data, in which the increase found in the younger ages over the last decade was higher among girls than among boys (16).

The geographical distribution of type 1 diabetes risk within the island was studied by mapping the relative risk at the municipality level. The standardized morbidity ratio observed in each small area varied largely from 0 to  $\geq 2$  and thus it gave no useful information. A Bayesian approach to the estimation of area-specific RR (Fig. 2A) showed that the highest risk areas were located in the southern and eastern parts of the island, whereas the lowest risk areas were in the northeastern part. However, the map of PP (Fig. 2B) shows that the RR was homogeneously distributed over the island with the exception of the Sassari municipality,

where the RR is  $< 0.95$ . A value of PP within the range of 0.25–0.75 indicates that the data do not provide any evidence of an RR being higher or lower than 1.


Furthermore, by comparing the maps of RR in 1989–1994 and in 1995–1999,

the geographical distribution of type 1 diabetes risk seemed almost unchanged and it remained homogeneously distributed (data not shown). However, the RR tended to be higher in the Campidano plain (the largest plain in Sardinia, located

**Table 2**—Time trend of type 1 diabetes incidence in children  $< 15$  years of age in Sardinia 1989–1999

Sex and age-group	Incidence rate ratio	95% CI	Percent increase	P
Boys				
0–4 years	1.043	0.991–1.097	4.3	0.104
5–9 years	1.006	0.967–1.047	0.6	0.755
10–14 years	1.036	1.001–1.071	3.6	0.041
0–14 years*	1.027	1.004–1.051	2.7	0.021
Girls				
0–4 years	1.038	0.976–1.104	3.8	0.234
5–9 years	1.018	0.973–1.065	1.8	0.445
10–14 years	1.036	0.991–1.084	3.6	0.120
0–14 years*	1.029	1.001–1.059	2.9	0.045
Boys and Girls				
0–4 years†	1.041	1.001–1.082	4.1	0.044
5–9 years†	1.011	0.982–1.042	1.1	0.462
10–14 years†	1.036	1.008–1.064	3.6	0.010
0–14 years*†	1.028	1.010–1.047	2.8	0.002

Results of the Poisson regression analysis. \*Age-standardized incidence rate; †sex-standardized incidence rate.



**Figure 2**—Estimates of type 1 diabetes risk among children aged 0–14 years in the 375 Sardinian municipalities during 1989–1999. A: Bayesian smoothed estimates of RR. B: Bayesian PP that the risk in each municipality exceeds the overall incidence.

in the southwestern part of the island) in the latter period (1995–1999).

**CONCLUSIONS** — Sardinia is known to have a very high incidence of type 1 diabetes (second after Finland [8] and five to seven times higher than continental Italy [17]). The present work confirms once again the high incidence of type 1 diabetes in the island.

We found one of the highest male-to-female incidence ratios ever reported in the 0- to 14-year age-group (1.4). The highest male-to-female ratio was found in the oldest age-group, even if a trend of male-to-female ratio with increasing age was not identified. Some authors describe a weak association between high incidence and high male-to-female ratio (18,19). Sardinia seems to confirm this observation. A possible explanation of this phenomenon could be found in a re-

cent survey on type 1 diabetic patients from the U.S., the U.K., and Sardinia in which a male sex bias was described in patients carrying neither DR3 nor DR4 haplotypes, and this effect was more pronounced among Sardinian patients (20), 87% of whom are DR3 positive (21).

The results of the present study, characterized by a prospective registration of cases, confirmed that in Sardinia there is an increasing risk of type 1 diabetes, as already shown by the Sardinian military conscript data among boys over the last decades (4), and this increase is even higher in recent years. Similar increasing rates have been reported elsewhere, in several countries (2–6,22) and in Europe as a whole (8).

In the latter study (EURODIAB study), no increase of incidence was reported for Sardinia. The different results we obtained are attributable to the differ-

ent time intervals considered (1989–1998 in the EURODIAB study and 1989–1999 in the present study) and mostly to a different completeness of ascertainment between the two studies. In fact, the completeness of the Sardinian data was 87% in the EURODIAB study and 91% in the present study. The difference was even higher if we consider the last years. This might be due to the fact that the new cases added to the present analysis were diagnosed in 1997–1998, which led to a higher incidence in the last years compared with the EURODIAB study and contributed toward a statistically significant temporal trend.

The increase of incidence observed in our analysis could be due to a higher degree of ascertainment over the last years compared with the previous ones, but even this seems not to be the case. In fact, the completeness of ascertainment varies

from 91 to 97% between 1989 and 1993 down to 82–92% between 1994 and 1999. Thus, if we would have calculated the incidence rates using the number of cases corrected for the completeness of ascertainment, then we would have obtained an even higher increase than the one we actually reported.

The increase was the same for both sexes. It is of notice that in the Finnish study, with a longer period of observation, the increase was more pronounced among boys in the past whereas among girls only recently (16). The observation period of our study might be too short to lead to any significant difference in increase between sexes even if this could explain the high male-to-female ratio.

This increase is apparent also in the youngest age-group, even if this is not statistically different from the other age-groups. It is, however, to be noticed that the highest increase was found in this age-group (4.1%). This could be explained by too few cases in this age-group to reach statistical significance.

On the basis of this trend, we can extrapolate the incidence for the future years. The predicted incidence based on the linear trend obtained with the Poisson regression analysis will be 52.7/100,000 (95% CI 43.3–94.0) in 2010 and 69.5/100,000 (47.8–100.1) in 2020. If this trend would remain the same, type 1 diabetes will represent one of the heaviest burdens for the Sardinian health authorities.

Apparently, the geographical distribution of type 1 diabetes risk exhibited only a slight heterogeneity. The highest risk areas were located in the southern and eastern part of the island, and the lowest risk areas were in the northwestern part. These findings confirm our previous study even if the risk seems to have leveled off in the whole island, with only the Sassari area at significant lower risk. A similar picture was obtained also from the data on military conscript for the birth cohorts 1974–1979 (23). Thus, the risk seems to be uniform within the island, and the comparison between the maps obtained for the first 5 years (1989–1994) and for the last 6 years (1995–1999) confirmed this impression. However, the RR tended to be higher in the Campidano plain (the largest plain in Sardinia located in the southwestern part of the island) over the period 1995–1999. A similar approach applied to Finland showed that the high-risk areas remained

mainly the same between 1987 and 1996 (24).

The described geographical and temporal homogeneity of diabetes risk and the rapid increase of incidence in a genetically stable and homogeneous population such as Sardinians suggest the role of an environmental factor uniformly distributed in the island. This makes it almost impossible to point out any possible etiological factors by applying ecological analysis to the Sardinian population. Examples for that could be no effect toward type 1 diabetes risk of temperature, rainfall, nitrate content of drinking water, milk, and casein intakes and a weak association with malaria (25–27). Comparison with other populations is then recommended, although this kind of comparative study has been unable so far to pinpoint any possible factor that may explain the high incidence of type 1 diabetes in Sardinia. In the recent analysis published by the EURODIAB Group (28), type 1 diabetes incidence was strongly associated in Europe with indicators of national prosperity, even though Sardinia lies far from the regression line between type 1 diabetes incidence and gross domestic product, coffee consumption, latitude, or milk intake, suggesting that the high incidence of type 1 diabetes on the island could be only partially explained by these factors.

The increase of incidence described here seems to be due to a widespread environmental agent operating in the genetically susceptible Sardinian population. However, these data are not able to explain the high incidence of the disease in Sardinia or its increase. We could speculate that some of the changes that occurred in the environment and lifestyle in Sardinia after World War II might have played a significant role. Transportation and connection within the island and to mainland Italy improved gradually after the 1950s as the social, economic, and hygiene standards improved. The environment also changed, particularly after the Rockefeller campaign that took place in the 1950s and was aimed at the eradication of malaria on the island. During that campaign, large amounts of DDT were spread all over the island, and its metabolites continue to be detected in the food chain. According to the studies by our group, the areas with past high malaria morbidity seem to be currently at lower risk of type 1 diabetes (27), leading

us to hypothesize that the *Plasmodium falciparum* might have had a protective role on type 1 diabetes, that it operated by selecting high-risk genes, or that it can be considered as an indicator of other parasitic infection that could be protective against type 1 diabetes. The disappearance of malaria might be then an epiphenomenon of some radical changes occurring in the Sardinian environment during and after the 1950s. This is in accordance with the so-called “hygiene hypothesis” (29,30), according to which the slow but continuous decline of infestation by intestinal helminthes and by many other ectoparasites and the reduction of morbidity and mortality of infectious disease brought about by changing lifestyles has resulted in disorders of the immune control and increases of immunomediated diseases. Animal data and epidemiological deductions are in accordance with this hypothesis. The high reduction of microbial exposure in infancy occurred in the last decade also in Sardinia, and the improved social standards might be responsible for the increase of several diseases in a genetically susceptible population. In fact, in Sardinia the almost complete disappearance of parasitic diseases and the eradication of polio, malaria, and tuberculosis have coincided with an increase of atopic and immunomediated diseases (type 1 diabetes, multiple sclerosis, Crohn’s disease, celiac disease). Unfortunately, very few data, if any, are available to support this hypothesis in Sardinia.

Furthermore, other changes occurred in the last 50 years in Sardinia (for example, there is better access to health care, high-caloric foods, fat, and meat proteins); the weight and height of the population also increased. Several reports linked linear growth and weight gain to the increased incidence of type 1 diabetes (31,32). Unfortunately, once again, no data are available on the growth velocity or the increase of obesity among Sardinian children in the past decades, even though it seems that the prevalence of overweight in Sardinian schoolchildren (12–24%) is comparable with the prevalence from mainland Italy (10–32%) (33).

In conclusion, the high incidence of type 1 diabetes is still increasing in Sardinia, confirming the increasing trend of prevalence among young boys shown by the analysis on the conscript registry (23).

In fact, our group already studied the temporal trend of type 1 diabetes point prevalence among young men aged 18 years at the moment of the conscript examination. This survey gave an increasing trend with the highest point prevalence of 4.92/1,000 registered in the last birth cohort of 1979. The Sardinian type 1 diabetes registry gave a cumulative incidence for young men aged 18 years of 6/1,000, which was higher than the point prevalence estimated from the military registry (23). This figure leads us to hypothesize a continuous, still ongoing increase of type 1 diabetes risk in Sardinian children and adolescents, at least among boys. Further investigations are needed to clarify whether this increase is mainly due to a peculiar genetic background, an environmental factor, or both. Given the homogeneous distribution of type 1 diabetes risk within the island, these studies should compare the Sardinian population with populations at different levels of risk.

## APPENDIX

### Members of the Sardinian Type 1 Diabetes Epidemiology Study Group

Coordinator: M. Songini; Secretariat Assistance: M. Porceddu; Coordinating Center: A. Casu, V.M. Cambuli; Study Group: E. Angius, G. Baule, L. Bellu, P. Bulciolu, A. Cabras, S. Cabras, L. Carboni, L. Carreras, M.A. Chessa, A.M. Cicalò, R. Cirillo, C. Clausi, M.T. Contigiani, A. Corda, E. Cossu, G. Delitala, L. De Luna, A. Dessupoiu, S. Erre, G. Fanciulli, A. Farris, A. Filigheddu, G.D. Filigheddu, O. Frongia; P. Frongia, T. Frulio, Z. Gambula, A. Gigante, M. Isoni, G. Idda, N. Landis, G. Lecca, S. Ledda, M. Maioli, M.C. Mannazzu, A.M. Marinaro, U. Marongiu, A. Masia, A. Massidda, F. Mastinu, G. Melis, L. Mereu, A.F. Milia, B. Nieddu, A. Ogana, P. Olmeo, A. Pacifico, G. Pala, C. Pipitone, G. Piredda, G. Pisano, R. Ricciardi, C. Ripoli, F. Sanciu, E. Secchi, V. Sica, M. Soro, A. Sturnara, P. Tronci, F. Velluzzi, and M.A. Zedda.

**Acknowledgments**—The authors thank Prof. F. Carle and Dr. R. Gesuita from the RIDI Study Group (Italian Registry for Insulin-Dependent Diabetes) for their comments and suggestions.

## References

- Gale EAM: The rise of childhood type 1 diabetes in the 20th century. *Diabetes* 51: 3353–3361, 2002
- Diabetes Epidemiology Research International Group: Secular trends in incidence of childhood IDDM in 10 countries. *Diabetes* 39:858–864, 1990
- Felterbower RG, McKinney PA, Parslow RC, Stephenson CR, Bodansky HJ: Type 1 diabetes in Yorkshire, U.K.: time trends in 0–14 and 15–29-years-old, age at onset and age-period-cohort modeling. *Diabet Med* 20:437–441, 2003
- Bruno G, Merletti F, Biggeri A, Cerutti F, Grosso N, De Salvia A, Vitali E, Pagano G, Piedmont Study Group for Diabetes Epidemiology: Increasing trend of type 1 diabetes in children and young adults in the province of Turin (Italy): analysis of age, period and birth cohort effects from 1984 to 1996. *Diabetologia* 44:22–25, 2001
- Gyurus E, Green A, Patterson CC, Soltesz G, the Hungarian Childhood Diabetes Epidemiology Study Group: Dynamic changes in the trends in incidence of type 1 diabetes in children in Hungary (1978–98). *Pediatr Diabetes* 3:194–199, 2002
- Schober E, Schnider U, Waldhor T, Tuomilehto J, Austrian Diabetes Incidence Study Group: Increasing incidence of IDDM in Austrian children: a nationwide study 1979–1993. *Diabetes Care* 18: 1280–1283, 1995
- Onkamo P, Vaananen S, Karvonen M, Tuomilehto J: Worldwide increase in incidence of type 1 diabetes: the analysis of the data in published incidence trends. *Diabetologia* 42:1395–1403, 1999
- Green A, Patterson CC, on behalf of the EURODIAB TIGER Study Group: Trends in the incidence of childhood-onset diabetes in Europe 1989–1998. *Diabetologia* 44 (Suppl. 3):B3–B8, 2001
- Songini M, Loche M, Muntoni Sa, Stabellini S, Coppola A, Dessì G, Green A, Bottazzo GF, Muntoni SE: Increasing prevalence of juvenile type 1 (insulin-dependent) diabetes mellitus in Sardinia: the military service approach. *Diabetologia* 36:547–552, 1993
- Cavalli-Sforza LL, Menozzi P, Piazza A: Sardinia. In *The History and Geography of Human Genes*. Cavalli-Sforza LL, Menozzi P, Piazza A, Eds. Princeton, NJ, Princeton University Press, 1994, p. 512–517
- Songini M, Bernardinelli L, Clayton D, Montomoli C, Pascutto C, Ghislandi M, Fadda D, Bottazzo GF, the Sardinian IDDM Study Groups: The Sardinian IDDM Study: 1. Epidemiology and geographical distribution of IDDM in Sardinia during 1989 to 1994. *Diabetologia* 41: 221–227, 1998
- Green A, Gale EAM, Patterson CC: Incidence of childhood-onset insulin-dependent diabetes mellitus: the EURODIAB ACE Study. *Lancet* 339:905–909, 1992
- ISTAT (Istituto Nazionale di Statistica) popolazione residente al 1 gennaio [article online], 2001. Available from <http://www.demo.istat.it/pop1/start.html>. Accessed 15 April 2002
- Bruno G, LaPorte R, Merletti F, Biggeri A, McCarty D, Pagano G: National diabetes programs: application of capture-recapture to “count” diabetes? *Diabetes Care* 17: 548–556, 1994
- Bernardinelli L, Clayton D, Pascutto C, Montomoli C, Ghislandi M, Songini M: Bayesian analysis of space time variation in disease risk. *Stat Med* 14:2433–2443, 1995
- Karvonen, Pitkaniemi J, Tuomilehto J, the Finnish Childhood Diabetes Registry Group: The onset age of type 1 diabetes in Finnish children has become younger. *Diabetes Care* 22:1066–1070, 1999
- Cherubini V, Chiarelli F, Altobelli E, Verrotti A, Carle F: Regional variability in the epidemiology of childhood diabetes in Italy. *J Pediatr Endocrinol Metab* 10:471–478, 1997
- Gale EAM, Gillespie KM: Diabetes and gender. *Diabetologia* 44:3–15, 2001
- Karvonen M, Pitkaniemi M, Pitkaniemi J, Kohtamäki K, Tajima N, Tuomilehto J: Sex difference in the incidence of insulin-dependent diabetes mellitus: an analysis of the recent epidemiological data: World Health Organization DIAMOND Project Group. *Diabetes Metab Rev* 13:275–291, 1997
- Cucca F, Goy JV, Kawaguchi Y, Esposito L, Merriman ME, Wilson AJ, Cordell HJ, Bain SC, Todd JA: A male-female bias in type 1 diabetes and linkage to chromosome Xp in MHC HLA-DR3-positive patients. *Nat Genet* 19:301–302, 1998
- Cucca F, Muntoni F, Lampis R, Frau F, Argiolas L, Silvetti M, Angius E, Cao A, De Virgili S, Congia M: Combinations of specific DRB1, DQA1, DQB1 haplotypes are associated with insulin-dependent diabetes mellitus in Sardinia. *Hum Immunol* 37:85–94, 1993
- Tuomilehto J, Karvonen M, Pitkaniemi J, Virtala E, Kohtamäki K, Toivanen L, Tuomilehto-Wolf E, the Finnish Childhood Type 1 Diabetes Registry Group: Record high incidence of type I (insulin-dependent) diabetes mellitus in Finnish children. *Diabetologia* 42:655–660, 1999
- Casu A, Pascutto C, Bernardinelli L, Songini M: Bayesian approach to study the temporal trend and the geographical variation in the risk of type 1 diabetes: the Sardinian Conscript Type 1 Diabetes Registry. *Pediatr Diabetes* 5:32–38, 2004
- Rytönen M, Ranta J, Tuomilehto J, Karvonen M, for the SPAT Study Group and

- the Finnish Childhood Diabetes Registry Group: Bayesian analysis of geographical variation in the incidence of type 1 diabetes in Finland. *Diabetologia* 44 (Suppl. 3): B37–B44, 2001
25. Casu A, Carlini M, Contu A, Bottazzo GF, Songini M: Type 1 diabetes in Sardinia is not linked to nitrate levels in drinking water. *Diabetes Care* 23:1043–1044, 2000
  26. Casu A, Fadda M, Bottazzo GF, Elliot RB, Songini M: Type 1 diabetes and cow milk in Sardinia: casein variant consumption (Abstract). *J Pediatr Endocrinol Metab* 14 (Suppl. 3):1018, OP-5, 2001
  27. Bernardinelli L, Pascutto C, Montomoli C, Komakec J, Gilks W, Songini M, Fiorani O, Lisa A, Zei G, Solinas G, Bottazzo GF: Bayesian analysis of ecological data for studying the association between insulin-dependent diabetes mellitus and malaria. In *Statistics for the Environment 4: Pollution Assessment and Control*. Barnett V, Stain A, Feridun Turkman K, Eds. New York, John Wiley & Sons, 1999, p. 29–47
  28. Patterson CC, Dahlquist G, Soltész G, Green A, the EURODIAB ACE Study Group: Is childhood-onset type I diabetes a wealth-related disease: an ecological analysis of European incidence rates. *Diabetologia* 44 (Suppl. 3): B9–B16, 2001
  29. Gale EA: A missing link in the hygiene hypothesis? *Diabetologia* 45:588–594, 2002
  30. Palmas C, Gabriele F, Conchedda M, Bor-toletti G, Ecça AR: Causality or coincidence: may the slow disappearance of helminths be responsible for imbalances in immune control mechanisms? *J Helminthol* 77:147–153, 2003
  31. Wilkin TJ: The accelerator hypothesis: weight gain as the missing link between type 1 and type 2 diabetes. *Diabetologia* 44:914–922, 2001
  32. Hypponen E, Virtanen SM, Kenward MG, Knip M, Akerblom HK, Childhood Diabetes in Finland Study Group: Obesity, increased linear growth, and risk of type 1 diabetes in children. *Diabetes Care* 23: 1755–1760, 2000
  33. Velluzzi F, Di Liberto M, Secci G, Calia MA, Mastinu R, Pilleri A, Cocco PL, Balestrieri A, Mariotti S, Loviselli A: Prevalenza del sovrappeso e dell'obesità nella scuola media inferiore in Sardegna (Abstract). *Riunione Sci Soc Ital Obes* A21, 2003